

<b>PVAS: Paediatric Vasculitis Activity Score</b>		Center/Physician:	
		Training Case/Patient No:	
<p>ÿ Tick "Active" box <b>only</b> if abnormality due to active vasculitis is newly present or worse over the last 4 weeks or persists for less than 3 months. After that, if ALL items are persistent and represent smouldering/low grade/grumbling disease, and there are no new/worse features, please tick the box at the bottom right corner. At the very first assessment all active items are considered as new/worse. If there are no abnormalities in a system, please tick the "None" box. For items present longer than 3 months refer to the Paediatric Vasculitis Damage Index (PVDI) to score damage.</p>			
		None	Active
<b>1. General</b>			
Myalgia			
Arthralgia or arthritis			
Fever $\geq 38.0^{\circ}\text{C}$			
Weight Loss $\geq 5\%$ body weight			
<b>2. Cutaneous</b>			
Polymorphous exanthema			
Livedo			
Panniculitis			
Purpura			
Skin nodules			
Infarct (nail edge lesion, splinter haemorrhage)			
Ulcer (full-thickness necrosis)			
Gangrene (extensive necrosis)			
Other skin vasculitis (specify below)			
<b>3. Mucous membranes/eyes</b>			
Mouth ulcers/granulomata			
Genital ulcers			
Adnexal inflammation			
Significant proptosis			
Red eye (Epi)scleritis			
Red eye conjunctivitis/ blepharitis/keratitis			
Uveitis			
Blurred vision			
Sudden visual loss			
Retinal vasculitis/retinal vessel thrombosis/retinal exudates/haemorrhages			
<b>4. ENT</b>			
Nasal discharge/crusts/ulcers/granuloma			
Paranasal sinus involvement			
Subglottic stenosis/ hoarseness /stridor			
Conductive hearing loss			
Sensorineural hearing loss			
<b>5. Chest</b>			
Wheeze or expiratory dyspnea			
Endobronchial/endotracheal involvement			
Nodules or cavities			
Pleural effusion/pleurisy			
Infiltrate			
Massive haemoptysis/Alveolar haemorrhage			
Respiratory failure			
<b>6. Cardiovascular</b>			
Loss of pulses			
Bruits over accessible arteries			
Blood pressure discrepancy			
Caudication of extremities			
Ischaemic cardiac pain			
Cardiomyopathy			
Congestive cardiac failure			
Valvular heart disease			
Pericarditis			
<b>7. Abdominal</b>			
Abdominal pain			
Peritonitis			
Blood in stools or bloody diarrhoea			
Bowel ischaemia			
<b>8. Renal</b>			
Hypertension >95th centile (for height)			
Proteinuria >0.3 g/24h, >20mg/mmol creatinine			
Haematuria $\geq 2+$ or 5 rbc/hpf or red cell casts			
GFR 50-80ml/min/1.73 m <sup>2</sup>			
GFR 15-49 ml/min/1.73 m <sup>2</sup>			
GFR <15 ml/min/1.73m <sup>2</sup>			
Rise in creatinine > 10% or			
Creatinine clearance (GFR) fall > 25%			
<b>9. Nervous system</b>			
Headache			
Meningitis/encephalitis			
Organic confusion/cognitive dysfunction			
Seizures (not hypertensive)			
Stroke			
Cord lesion			
Cranial nerve palsy			
Sensory peripheral neuropathy			
Motor mononeuritis multiplex			
<b>10. OTHER</b>			
<b>PERSISTENT DISEASE ONLY :</b>			
Tick here if there is no new/worse abnormality present in ANY of the systems above and active items represent persistently active disease			ÿ